

PROSTALGIN

TRADITIONAL INDICATIONS

1. Prostate Enlargement (Benign Prostatic Hyperplasia)
2. Urinary Calculi
3. Eczema
4. Dermatitis and allergic skin conditions
5. Repeated Epididymitis infections
6. Chronic epididymitis



DOSAGE and DIRECTIONS



For Adults 18 yrs or above, two (2) Capsules twice daily (every 12 hours) in empty stomach with lukewarm water.

Please give minimum 30 min to take any food after taking PROSTALGIN.

Take optimum water to enhance elimination of released toxins from the body.

INGREDIENTS

Supplement Facts		
Serving Size: 2 Capsule (500 mg each)		
Servings per container: 60		
Each Serving contains		%DV
Andrographis Paniculata	100 mg.	*

177 Alpha Ave, Muskegon, MI 49441-2659, United States of America.
Web: www.gurusgarden.com

Curcuma Longa	100 mg.	*
Commiphora Mukul	90 mg.	*
Plumbago Zeylanica	25 mg.	*
Embelia Ribes	25 mg.	*
Coriandrum Sativum	25 mg.	*
Cyperus Rotundus	25 mg.	*
Tinospora Cordifolia	20 mg.	*
* Daily value (DV) not established.		

HEALING CRISIS

There is no Healing crisis.